

3-6-2020

Metastatic Spinal Cord Compression Mimicking Cauda Equina Syndrome Illustrating Importance of Adequate Diagnostic Imaging

Jonathan L. Holt

Joseph Hill

HCA Healthcare, Joseph.Hill2@hcahealthcare.com

Farzin Farhandnejad

HCA Healthcare, farzin.farhangnejad@hcahealthcare.com

Karyn Doddy

HCA Healthcare

Se Won Lee

HCA Healthcare, sewon.lee@hcahealthcare.com

Follow this and additional works at: <https://scholarlycommons.hcahealthcare.com/pmr>



Part of the [Neurology Commons](#), [Pathological Conditions, Signs and Symptoms Commons](#), and the [Rehabilitation and Therapy Commons](#)

Recommended Citation

Holt JL, Hill J, Farhangnejad F, Doddy K, Lee SW. Metastatic Spinal Cord Compression Mimicking Cauda Equina Syndrome Illustrating Importance of Adequate Diagnostic Imaging. Poster presented at: Association of Academic Physiatrists Meeting; March 4-9, 2020; Orlando, FL.

This Poster is brought to you for free and open access by the Research & Publications at Scholarly Commons. It has been accepted for inclusion in Physical Medicine & Rehabilitation by an authorized administrator of Scholarly Commons. For more information, please contact lindsay.million@hcahealthcare.com.

Metastatic Spinal Cord Compression Mimicking Cauda Equina Syndrome Illustrating Importance of Adequate Diagnostic Imaging

Jonathan L. Holt, MSIV¹, Joseph Hill, D.O.², Farzin Farhang-Nejad, M.D.², Karyn Doddy, M.D.², Se Won Lee, M.D.²

¹University of Nevada Reno School of Medicine, ²Department of Physical Medicine and Rehabilitation/ Sunrise Health Graduate Medical Education Consortium, Mountain View Medical Center

Introduction

MRI is the gold standard for diagnosis of spinal cord compression (SCC) with 93% sensitivity and 97% specificity.

- Inability to obtain the MRI can hinder proper management in the emergent case of SCC.
- CT Myelography is a valuable imaging tool for SCC patients unfit for MRI, and should be the next step among those who show "red flag" signs of pain, motor/sensory deficits, and/or autonomic deficits.
- Localizing the level of spinal cord compromise, is paramount to managing SCC as it is a medical emergency, and may be reversible if it is identified early and treatment is started quickly.

Case Summary

History - Patient is a 66-year-old morbidly obese male with a recent diagnosis of prostatic cancer and mixed bony metastasis of the axial skeleton (Fig 1). Initial attempts to obtain MRI after the diagnosis were failed due to the patient's large body habitus. Two prior falls 27 and 16 days before the current admission resulted in a right humeral compression fracture and left sacral ala fracture respectively. The mechanism of both falls was reported to be sudden onset weakness in the lower extremities. CT lumbar spine and pelvis on current admission was unchanged: diffuse metastatic disease throughout the spine, severe spinal stenosis L2-L5, no definitive epidural tumor, and left sacral ala fracture. Current presentation is an acute onset of low back pain, bilateral lower extremity weakness, inability to ambulate, paresthesia, and one recent episode of urinary incontinence leading to patient's admission by Neurosurgery for suspected Cauda Equina Syndrome (CES). Corticosteroids were started and emergent L2-L5 laminectomy was performed. Progress notes documented improved motor strength and sensation in the lower extremities immediately following lumbar decompression. PM&R was consulted and admitted the patient for rehabilitation of a suspected cauda equina syndrome 2 days after the lumbar laminectomy.

Physical Exam at time of Inpatient Rehab admission

- Bilaterally upper ext strength: 5/5. Limited use of right shoulder due to previous proximal humerus fracture.
- Lower ext strength: Left ankle dorsiflexion 2/5. Right knee flexion 4/5 and Left knee flexion 3/5. Hip flex bilat 2/5.
- CNII-XII grossly intact, 1+ Reflexes, 5 beats of clonus in bilateral ankles
- Presented to rehab with urinary retention that was eventually treated with suprapubic catheter.

Rehabilitation – The patient did not functionally progress during his inpatient rehabilitation due to pain and limited mobility. During the 7th day the patient developed acute worsening of paralysis and complete paresthesia of his lower extremities. He was readmitted to the ICU and a spinal myelogram (Fig. 2) was performed revealing a contrast block at T6-T7 consistent with an epidural tumor. A T4-T8 laminectomy and epidural tumor evacuation was subsequently performed by neurosurgery.

- Outcome - Ten days after the second spinal decompression/evacuation operation, the patient was readmitted to inpatient rehabilitation with no lower extremity motor function and decreased sensation below the T8 dermatome. He was eventually discharged to SNF on hospice.

Images




Fig. 1: CT of lumbosacral spine




Fig. 2: CT myelogram of thoracic spine

Discussion

Although this patient likely received appropriate surgical intervention that was correlated to the CT imaging, symptoms, and history of prostate cancer, the delay in obtaining full spinal imaging with CT myelogram and diagnosis of a second compressive epidural tumor is a non-dismissable factor that may have lead to a less favorable outcome for this patient.

Initial treatment with steroids may have halted or decreased the inflammation caused by the epidural tumor in the thoracic spine which may have contributed to the initial improvement in symptoms following the first lumbar spine decompressive surgery.

If CT myelogram was obtained at admission and decompressive surgeries were performed at lumbar and thoracic levels, this patient would be left with paresis in his legs instead of plegia in legs in abdominal core yielding a better prognosis.

Conclusion

MRI is the gold standard for diagnosis of spinal cord compression and cauda equina syndrome in the setting of metastatic cancer.

Complete spinal imaging should be obtained if there are concerns for urgent SCC and imminent neurological damage.

- If MRI is unobtainable, CT myelogram should be the next step to diagnose suspected MSCC.

There are statistically significant higher ambulatory rates in patients receiving high-dose dexamethasone before radiotherapy

Following confirmation of diagnosis CES, start corticosteroids and take the patient to surgery for emergent decompression.

Following confirmation of the diagnosis of SCC, start corticosteroids and take the patient to surgery for decompression/stabilization and follow up with radiation therapy.

References

- Boussios S, Cooke D, Hayward C, et al. Metastatic Spinal Cord Compression: Unraveling the Diagnostic and Therapeutic Challenges. *Anticancer Research*. 2018;38(9):4987-4997. doi:10.21873/anticancer.12817.
- Husband DJ, Grant KA, Romaniuk CS. MRI in the diagnosis and treatment of suspected malignant spinal cord compression. *The British Journal of Radiology*. 2001;74(877):15-23. doi:10.1259/bjr.74.877.740015.
- Loblaw DA, Perry J, Chambers A, Laperriere NJ. Systematic Review of the Diagnosis and Management of Malignant Extradural Spinal Cord Compression: The Cancer Care Ontario Practice Guidelines Initiative's Neuro-Oncology Disease Site Group. *Journal of Clinical Oncology*. 2005;23(9):2028-2037. doi:10.1200/jco.2005.00.067.
- Long B, Koyfman A, Gottlieb M. Evaluation and management of cauda equina syndrome in the emergency department. *The American Journal of Emergency Medicine*. 2020;38(1):143-148. doi:10.1016/j.ajem.2019.158402.
- Mcquail PM, Mccartney BS, Baker JF, Jaadan M, McCabe JP. Management of Metastatic Spinal Cord Compression in Ireland: Are Surgeons Overlooked? *International Journal of Spine Surgery*. August 2018:5051. doi:10.14444/5051.
- Ohashi M, Hirano T, Watanabe K, et al. Preoperative prediction for regaining ambulatory ability in paretic non-ambulatory patients with metastatic spinal cord compression. *Spinal Cord*. 2016;55(5):447-453. doi:10.1038/sc.2016.145.