Case Report

Jacuzzi-Induced Pneumoperitoneum: Case Report and Literature Review

Obteene Azimi-Ghomi, MD,¹ Vasiliy Ovakimyan, MD,¹
Dieter Brummond, MD,¹ Gerardo Kahane, MD¹

Abstract

Description

Radiographically-detected pneumoperitoneum in the setting of acute onset abdominal pain is a classic case among surgical emergencies. The spectrum of etiology ranges from benign to catastrophic, and the ability to differentiate depends in large part on the history and physical exam findings. Included among non-surgical types of pneumoperitoneum are (1) post-operative, (2) diagnostic/experimental, (3) gynecologic tract air entry, (4) the difference in intra- and extra-alveolar pressures, pneumothorax-associated and (5) miscellaneous. Jacuzzi-induced pneumoperitoneum is an extremely rare form of non-surgical pneumoperitoneum. We discuss a case of a 32-year-old female presenting with pneumoperitoneum and acute onset abdominal pain secondary to vaginal insertion onto a Jacuzzi jet. We describe the workup and management of this case and perform a literature review of reported cases of Jacuzzi-induced pneumoperitoneum. Although its presentation is dramatic, a prudent physical exam, history and appropriate imaging are key to the management of this non-surgical condition.

Keywords

pneumoperitoneum; pneumoperitoneum/etiology; pneumoperitoneum/diagnostic imaging; peritoneal diseases; Jacuzzis; accidental injuries

Introduction

Radiographically-detected pneumoperitoneum in the setting of acute onset abdominal pain is a classic case among surgical emergencies. The spectrum of etiology ranges from benign to catastrophic, and the ability to differentiate depends in large part on the history and physical exam findings. Included among non-surgical types of pneumoperitoneum are (1) post-operative, (2) diagnostic/experimental, (3) gynecologic tract air entry, (4) the difference in intra- and extra-alveolar pressures, pneumothorax-associated and (5) miscellaneous.¹⁻⁵ Jacuzzi-induced pneumoperitoneum is an extremely rare form of non-surgical pneumoperitoneum. Although its presentation is dramatic, a prudent physical exam, history and appropriate imaging are key to the management of this non-surgical condition.

Case Presentation

A 32-year-old female with no past medical history presented with acute onset abdominal pain localized to the epigastrium. Vital signs were normal, and the patient was afebrile. Laboratories were normal, demonstrating hemoglobin of 13.6 g/dL, white blood cell count of 8000 and lactic acid of 0.8 mm. Upon physical exam, the patient's abdomen was peritonitic. The patient underwent a contrast-enhanced computed tomography (CT) of the abdomen and pelvis revealed pneumoperitoneum localized to the right upper quadrant with mild gastric and small bowel mucosal thickening, raising suspicion for a perforated duodenal ulcer. (**Figures 1 and 2**)

Emergency exploration was indicated, and the patient's clinical stability permitted a laparo-

Author affiliations are listed at the end of this article.

Correspondence to:
Obteene Azimi-Ghomi, MD
General Surgery
Department
Aventura Hospital and
Medical Center
20900 Biscayne Blvd.
Aventura, Florida, 33180
(Obteene.azimighomi@hcahealthcare.com)

Figure 1. Coronal view of CT abdomen/pelvis demonstrating pneumoperitoneum with multiple foci of free air under the diaphragm and around the liver (arrows).

scopic approach. Laparoscopic examination of the stomach, duodenum and small and large bowel failed to identify a perforation. An intra-operative esophagogastroduodenoscopy also failed to demonstrate any perforations or luminal irregularity. The decision was made to leave a drain and transfer the patient to the ICU with nasogastric decompression and antibiotic coverage. Post-operatively, discussion with the patient's spouse elucidated the fact that prior to arrival to our facility, they had been engaging in sexual activity in their Jacuzzi, with the patient having inserted the Jacuzzi jets into her vagina for sexual pleasure. Further review of the imaging following this new knowledge demonstrated extensive air within the uterus, which had initially gone unnoticed.

(**Figure 3**) In light of this new information, the patient was observed for 24 hours with labs and vitals remaining stable. The abdominal exam remained unchanged. The patient was started on a liquid diet post-operative on day 1. The patient tolerated a normal diet and was discharged on postoperative day 2 in stable condition.

Discussion

Pneumoperitoneum is classically a surgical emergency, with perforated hollow viscera accounting for about 41–90% of cases.^{1,2} Perforated peptic ulcer disease is the most common surgical etiology to present with pneumoperitoneum, with 72% of cases demonstrating this finding.² Chest or abdominal plain film

Figure 2. Sagittal view of CT abdomen/pelvis demonstrating extensive free air throughout the abdomen (yellow arrows) and air within the uterus (red arrow).

Figure 3. Axial view of CT abdomen/pelvis demonstrating air within the fallopian tubes (arrows). Seen on “lung window.”

radiography is the primary diagnostic tool for detecting pneumoperitoneum.³ Computed tomography (CT) examinations, however, have been shown to be more sensitive than abdominal radiographs for the detection of free intraperitoneal air.³ Surgical causes of pneumoperitoneum typically require laparotomy for repair, resection or exclusion of the perforated viscus depending on the source.⁴ This fact is especially true in patients who are hemodynamically unstable or demonstrate laboratory abnormalities.⁴ Many reports describe pneumoperitoneum developing from non-surgical etiologies. Miller et. al in 1984 classified the causes of non-surgical pneumoperitoneum into 5 different etiologies: post-operative, diagnostic and experimental, gynecologic/fallopian tube air entry, difference in intra- and extra-alveolar pressures, pneumothorax-associated and miscellaneous.⁵

Multiple gynecologic causes of pneumoperitoneum have been described in the literature, including coitus,⁶⁻¹¹ oro-genital sex,¹²⁻¹⁵ post-partum knee-chest exercises,^{6,16,17} vaginal douching,^{6,7,21} pelvic inflammatory disease,⁵ hysterosalpingography,⁵ pelvic examinations,^{6,7,18-20} and aquatic sports.²² Jacuzzi-induced pneumoperitoneum is an extremely rare cause of pneumoperitoneum, with only three previously reported cases in the literature, which are described below.²³⁻²⁵

The first reported case in the literature was from the United Kingdom in 2000, involving

a 56-year-old female who presented with abdominal pain following Jacuzzi use hours earlier.²³ The patient was mildly tachycardic and peritonitic. Pneumoperitoneum was demonstrated on abdominal radiographs. An exploratory laparotomy was performed demonstrating clear fluid and gas within the abdomen without evidence of perforated viscus. The patient was treated with antibiotics and discharged 48 hours later.

The second reported case was from the United States in 2004 involving a 49-year-old female who developed abdominal pain one day after slipping and falling in a Jacuzzi tub where she reported a feeling of air and water rushing into her vagina.²⁴ The patient's vitals and labs were normal. A physical exam demonstrated peritonitis. A CT of the abdomen and pelvis demonstrated a considerable amount of free intraperitoneal gas and equivocal mural thickening of the second portion of the duodenum. The patient was admitted for observation and did not undergo surgical exploration. She underwent an upper gastrointestinal series with water-soluble contrast that did not demonstrate any abnormality. The patient was subsequently discharged home.

The third and most recently reported case was from Hungary in 2013 involving a 61-year-old female who presented with diffuse abdominal pain without any other symptoms.²⁵ Chest and abdominal radiographs demonstrated free air under the diaphragms. The patient underwent

surgical exploration with a laparotomy, and no evidence of a perforated viscus or a source of peritonitis was found. Post-operatively, the patient admitted to her complaints developing after using the Jacuzzi.²⁵

All 3 cases presented with severe abdominal pain following Jacuzzi use. The time frame for symptom development ranged between several hours to a day following Jacuzzi usage.²³⁻²⁵ All patients demonstrated peritonitis upon clinical examination with normal vital signs and laboratories. Two patients were diagnosed with abdominal radiography. The remaining patient underwent both radiography and CT imaging. The patients diagnosed with pneumoperitoneum solely by plain film radiography underwent exploratory laparotomy.^{23,25} The third patient was treated conservatively in a non-operative manner.²⁴ All patients were discharged home in stable condition within 72 hours of admission.

Conclusion

Jacuzzi-induced pneumoperitoneum is a rare form of non-surgical pneumoperitoneum. A detailed physical examination and medical history are integral to diagnosis. As with other types of pneumoperitoneum, chest or abdominal radiography can demonstrate air beneath the diaphragms. A CT of the abdomen and pelvis is the preferred imaging modality as it can characterize the extent of pneumoperitoneum, detect any obvious signs of viscus perforation and can examine the pelvic organs for any air that could indicate pelvic entry. An accurate history and physical examination along with CT imaging and stable vitals and laboratory values can help diagnose this phenomenon. Patients with the appropriate history of Jacuzzi usage in combination with normal vital signs, laboratories and findings of an isolated pneumoperitoneum on CT imaging can be safely observed. Empiric broad-spectrum antibiotics are recommended until the patient demonstrates persistent clinical stability with improved abdominal pain on examination. Should the patient require exploration, a diagnostic laparoscopic examination of the abdomen can help to rule out a perforated viscus and avoid any extensive surgical interventions.

Consent for Publication

The patient in the aforementioned case report

gave verbal and written consent for the case to be published.

Authors' Contributions

Obteene Azimi-Ghomi was involved in the sourcing of dates and resources for the article and the editing of images. Obteene Azimi-Ghomi, Vasiliy Ovakimyan, and Deiter Brummond were involved in the writing, editing, and formatting of the case report. Gerardo Kahane was involved in the development of the focus of the article and provided advice regarding contextualization and development of the article.

Acknowledgments

We would like to acknowledge the patient care team, including nursing, technician, radiology and operative support staff during the care of the patient.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

The authors are employees of Aventura Hospital and Medical Center, a hospital affiliated with the journal's publisher.

This research was supported (in whole or in part) by HCA Healthcare and/or an HCA Healthcare affiliated entity. The views expressed in this publication represent those of the author(s) and do not necessarily represent the official views of HCA Healthcare or any of its affiliated entities.

Author Affiliations

1. Aventura Hospital and Medical Center, Aventura, FL

References

1. Winek TG, Mosely HS, Grout G, Luallin D. Pneumoperitoneum and its association with ruptured abdominal viscus. *Arch Surg*. 1988;123(6):709-712. <https://doi.org/10.1001/archsurg.1988.01400300051008>
2. Kumar A, Muir MT, Cohn SM, Salhanick MA, Lankford DB, Katabathina VS. The etiology of pneumoperitoneum in the 21st century. *J Trauma Acute Care Surg*. 2012;73(3):542-548. <https://doi.org/10.1097/ta.0b013e31825c157f>

3. Pinto A, Miele V, Schillirò ML, et al. Spectrum of signs of pneumoperitoneum. *Semin Ultrasound CT MR*. 2016;37(1):3-9. <https://doi.org/10.1053/j.sult.2015.10.008>
4. Townsend C, Beauchamp RD, Evers BM, Mattox K. Acute abdomen. In: Townsend CM, ed. *Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice*. 20th ed. Elsevier; 2016:1120-1137.
5. Miller RE, Becker GJ, Slabaugh RD. Nonsurgical pneumoperitoneum. *Gastrointest Radiol*. 1981;6(1):73-74. <https://doi.org/10.1007/bf01890224>
6. Daly BD, Guthrie JA, Couse NF. Pneumoperitoneum without peritonitis. *Postgrad Med J*. 1991;67(793):999-1003. <https://doi.org/10.1136/pgmj.67.793.999>
7. Jacobs VR, Mundhenke C, Maass N, Hilpert F, Jonat W. Sexual activity as cause for non-surgical pneumoperitoneum. *JSLs*. 2000;4(4):297-300.
8. Im DD, Pak PS, Cua B, Feinberg E. Pneumoperitoneum after sexual assault in a patient who had hysterectomy 30 years ago: case report. *J Emerg Med*. 2012;42(5):540-542. <https://doi.org/10.1016/j.jemermed.2011.05.049>
9. Christiansen WC, Danzl DF, McGee HJ. Pneumoperitoneum following vaginal insufflation and coitus. *Ann Emerg Med*. 1980;9(9):480-482. [https://doi.org/10.1016/s0196-0644\(80\)80308-8](https://doi.org/10.1016/s0196-0644(80)80308-8)
10. Spaulding LB, Gallup DG. Pneumoperitoneum after hysterectomy. *JAMA*. 1979;241(8):825.
11. Tabrisky J, Mallin LP, Smith JA 3rd. Pneumoperitoneum after coitus. A complication due to uterine tube prolapse after vaginal hysterectomy. *Obstet Gynecol*. 1972;40(2):218-220.
12. Collins KA, Davis GJ, Lantz PE. An unusual case of maternal-fetal death due to vaginal insufflation of cocaine. *Am J Forensic Med Pathol*. 1994;15(4):335-339. <https://doi.org/10.1097/00000433-199412000-00010>
13. Fyke FE III, Kazmier FJ, Harms RW. Venous air embolism. Life-threatening complication of orogenital sex during pregnancy. *Am J Med*. 1985;78(2):333-336. [https://doi.org/10.1016/0002-9343\(85\)90445-0](https://doi.org/10.1016/0002-9343(85)90445-0)
14. Freeman RK. Pneumoperitoneum from oral-genital insufflation. *Obstet Gynecol*. 1970;36(1):162-164.
15. Varon J, Laufer MD, Sternbach GL. Recurrent pneumoperitoneum following vaginal insufflation. *Am J Emerg Med*. 1991;9(5):447-448. [https://doi.org/10.1016/0735-6757\(91\)90211-2](https://doi.org/10.1016/0735-6757(91)90211-2)
16. Dodek SM, Friedman JM. Spontaneous pneumoperitoneum. *Obstet Gynecol*. 1953;1(6):689-698.
17. Lozman H, Newman AJ. Spontaneous pneumoperitoneum occurring during postpartum exercises in the knee-chest position. *Am J Obstet Gynecol*. 1956;72(4):903-905. [https://doi.org/10.1016/0002-9378\(56\)90181-8](https://doi.org/10.1016/0002-9378(56)90181-8)
18. Mularski RA, Sippel JM, Osborne ML. Pneumoperitoneum: a review of nonsurgical causes. *Crit Care Med*. 2000;28(7):2638-2644. <https://doi.org/10.1097/00003246-200007000-00078>
19. Vlachou PA, Aslam M, Ntatsios A, Anagnostopoulos GK, Murphy P. Non-surgical pneumoperitoneum following cervical smear test. *Eur J Obstet Gynecol Reprod Biol*. 2007;131(2):240-241. <https://doi.org/10.1016/j.ejogrb.2006.04.025>
20. Cass LJ, Dow EC, Brooks JR. Pneumoperitoneum following pelvic examination. *Am J Gastroenterol*. 1966;45(3):209-211.
21. Walker MA. Pneumoperitoneum following a douche. *J Kans Med Soc*. 1942;43-55.
22. Johnson LB. A case report: Peritonitis after water-skiing. Fallbeskrivning: Peritonit efter vattenskidåkning. *Lakartidningen*. 1993;90(25):2383.
23. Inston N, Lake S. Pneumoperitoneum following Jacuzzi usage. *Ann R Coll Surg Engl*. 2000;82(5):350-351.
24. Williams TC, Kanne JP, Lalani TA. Jacuzzi jet-induced pneumoperitoneum. *Emerg Radiol*. 2004;10(5):259-261. <https://doi.org/10.1007/s10140-004-0330-y>
25. Farics A, Simon E, Parti K, Bátorfi J. Pneumoperitoneum after jacuzzi bathing -- an unusual cause for the development of intraperitoneal free air. *Jacuzzihasználatot követően kialakult pneumoperitoneum, avagy a szabad hasi levegő keletkezésének ritka esete. Magy Seb*. 2013;66(2):62-66. <https://doi.org/10.1556/maseb.66.2013.2.2>